

Caregiver reports of provider recommended frequency of blood glucose monitoring and actual testing frequency for youth with type 1 diabetes

Joyce P. Yi-Frazier and colleagues for the SEARCH for Diabetes in Youth Study Group

Authoritative, evidence-based recommendations suggest that children and adolescents with type 1 diabetes should monitor their blood glucose regularly and frequently (four times per day or more) but how feasible is this, socially and financially particularly in low resource societies or when families have to foot the bill themselves? This article is reproduced as published in *Diabetes Research and Clinical Practice (DRCP)* (2012) with the full list of authors, abstract, statistical methods, tables, acknowledgments and references available via the DRCP website. The setting is a multi-center study from the USA. Future issues of *Diabetes Voice* may explore these questions further in less well-resourced environments.

Frequent blood glucose monitoring – a cornerstone of diabetes management

Given the complexity and demanding nature of modern-day treatment regimens for type 1 diabetes, it is not surprising that reported adherence to treatment recommendations among youth with diabetes is low, particularly among adolescents.^{1,2} Understanding the factors that are related to adherence is critical. Studies have clearly demonstrated that persons with higher levels of adherence to their treatment regimens have better glycemic control.^{3,4} Since sub-optimal glycemic control is associated with an increased risk for the development of diabetes-related complications,⁵ youth with type 1 diabetes should strive to achieve and maintain good glycemic control, even at an early age. However, optimal glycemic control in youth is not often achieved, particularly by older youth.^{6,7}

Based on the findings of the Diabetes Control and Complications Trial,⁸ frequent blood glucose monitoring (BGM) has become a cornerstone of optimal diabetes management. Adherence to frequent BGM has been found to be an integral factor in achieving optimal glycemic control.^{1,9} Greater BGM has been reported to be associated with younger age and insulin pump use.^{9,10} In most cases, a linear increase in BGM per day resulted in improvements in glycated hemoglobin (A_{1c}),¹ but recent analyses have suggested that this effect may peak at 5 times/day.¹⁰

While previous research has shown that BGM is an important contributor to glycemic control, little is known about whether BGM by children with type 1 diabetes corresponds with the frequency of BGM recommended by their health care providers. Using data from

the SEARCH for Diabetes Study, these analyses explore demographic, family and clinical characteristics associated with provider recommended frequency of BGM, actual frequency of BGM, and concordance between these categories in children with type 1 diabetes as reported by child's caregiver. Additionally we explore the associations between recommended and reported frequency of BGM and A_{1c} measurements.

SEARCH for Diabetes in Youth is a multicenter study that conducts population-based ascertainment of youth with clinically diagnosed, non-gestational diabetes who are <20 at the time of diagnosis.¹¹ SEARCH has enrolled youth newly diagnosed with diabetes from 2002 through the present. Cases are ascertained from geographically defined populations in Ohio, Colorado, South Carolina and Washington, Indian Health Service beneficiaries from four American Indian populations, and enrollees in several health plans in California and Hawaii. Youth whose diabetes is not secondary to other conditions are invited to a SEARCH study visit. After obtaining informed consent and assent, physical measurements and fasting blood samples are collected from metabolically stable children, and questionnaires are administered. Youth whose diabetes was incident in 2002 through 2005 and who completed a baseline study visit were invited to return for follow-up visits.

The data included in this analysis are from children with type 1 diabetes who were 10 through 17 years of age and their parent/guardian ('caregivers') who accompanied them at their 24-month SEARCH follow-up visit where questions about their child's provider recommendation for frequency of BGM and their child's actual frequency of BGM

were asked. The response rate for this follow-up visit was 52%. Of the 390 participants who completed a follow-up visit, 94% (n = 385) completed the survey which included the questions about BGM.


Measures studied included demographic, clinical and metabolic variables.

Demographic variables included age at study visit, sex, race ethnicity, and insurance status. Insurance was categorized as private, state-funded (Medicaid/Medicare, etc.), other (which included student health clinics, military, Indian Health Services), or none. Family variables included family income which was categorized as <\$25K, \$25–49K, \$50–74K, \$75K+, do not know/refused, and family composition which was categorized as 2-parent household, 1-parent household, or other/unknown composition.

Clinical variables included duration of diabetes since diagnosis, insulin regimen, depression, and glycemic control. Duration of diabetes was the length of time between date of diagnosis and the 24-month visit. Insulin regimen was categorized as (1) basal-bolus using the insulin pump, (2) basal-bolus with glargine plus rapid-acting insulin, (3) multiple daily injections (MDI) with ≥3 injections/day, using glargine plus more than/ or other than rapid-acting insulin type, (4) MDI with ≥3 injections/day, using any insulin types excluding basal insulin, or (5) 1–2 injections/day, excluding glargine.⁶ Detemir and glulisine were not in clinical use during the data collection period. Depression was assessed based on the child's responses on the Center for Epidemiologic Studies Depression scale (CES-D).¹² This 20-item scale is a



Study participants are recently diagnosed young people with type 1 diabetes.



commonly used measure of depressive symptomatology in children age 10 years and older.^{13,4} For these analyses, we categorized the scores as minimally (0–15), mildly (16–23), and moderately/severely (24–60) depressed mood.^{15,16}

Glycemic control was assessed using blood samples shipped to a central laboratory (Northwest Lipid Research Laboratories, Seattle, WA) for analysis. An ion exchange unit (Variant II; Bio-Rad Diagnostics, Hercules, CA) quantified the glycated hemoglobin (A_{1c}) levels. Optimal age-specific goals for A_{1c} in children are <8.0% for 10–12 year olds, and <7.5% for 13–18 year olds.¹⁷

The frequency of blood glucose monitoring recommended by the child's provider as reported by the caregiver/parent – key data collected.

Caregivers of children with type 1 diabetes were asked to report the frequency of BGM recommended by their child's health care provider and the number of times per day their child conducted BGM over the last three months. Response options for both questions were 6 or more times daily, 4–5 times daily, 2–3 times daily, at least once daily, or do not know. Adherence to recommendations was determined by comparing their responses to questions about recommended and actual BGM frequency, and categorized as exceeded (child monitored more than recommended by provider), met (child monitored exactly as the provider recommended), or did not meet (child monitored less than what the provider recommended) their child's provider recommendation for daily BGM.

Detailed statistical analysis methods are given in the link as are the results tables.

Table 1 shows the demographic, family and clinical composition of the 390 eligible children with type 1 diabetes. They ranged from ages 10 to 17 (mean \pm SD = 14.0 \pm 2.3 years) with a mean diabetes duration of just over 3 years (37.4 months). About half were female (48%) and 74% were NHW. Youth 13–15 years had the highest mean A_{1c} values (9.2 \pm 1.9 for 13–15 year olds, 8.4 \pm 1.5 for 10–12 year olds, and 8.7 \pm 1.8 for 16–17 year olds; $p = .003$) and were least likely to be in adequate control based on their A_{1c} value (15% of 13–15 year olds, 41% of 10–12 year olds, and 27% 16+ met the goal; $p < .001$). Other demographic, clinical, and family characteristics did not differ significantly by age (data not shown).

A fifth of caregivers reported that their child's health care provider recommended blood glucose monitoring six times a day – every day. This advice varied considerably from centre

reported 2–3 times/day (Table 2, $n = 5$ with missing data). There was a significant difference in recommended frequency by study centers ($p = .04$, data not shown). Caregivers report of recommended frequency of BGM varied significantly by their child's age, race/ethnicity, household composition and family income as well as insulin regimen and glycemic control (Table 2). Younger, NHW participants, those from two parent households, those from families with higher incomes, and those on an insulin pump were significantly more likely to report being told to conduct BGM at least 6 times/day (Table 2). Additionally, caregivers of children who met their A_{1c} goal were more likely to report being told to check more frequently than those who did not meet their goal. The child's gender, health insurance status, duration of diabetes and depression scores were not associated with caregivers' report of providers' recommendations.

Of the caregivers surveyed, 26% reported that their child conducted BGM at least 6 times/day, 52% 4–5 times/day,

Higher family incomes and better glycaemic control were associated with more frequent blood glucose monitoring.

to centre and according to a number of characteristics of the person with diabetes and their families.

Of the caregivers surveyed, 22% reported that they were told by their health care provider their child should conduct BGM at least 6 times/day, 69% reported 4–5 times/day, and 8%

16% 2–3 times/day, and 6% reported fewer than 2 times/day (Table 2, $n = 2$ with missing data). There was no significant difference between participant responses by study center (data not shown). Caregivers of younger children, those from two parent households, those with higher incomes, those who were NHW, and those on an insulin

pump were more likely to report that their child conducted BGM at least 6 times/day (Table 2). Additionally, those children who met their A_{1c} goal were more likely to have caregivers who reported that they conduct BGM more frequently than those who did not meet their goal. There was no association between frequency of BGM and child's gender, health insurance status, duration of diabetes, or depressive symptoms.

A_{1c} was significantly lower for children whose caregivers reported that providers recommended BGM at least 6 times/day compared to those who reported that their providers recommended BGM 4–5 times/day, or fewer. A_{1c} values were also significantly lower for children who reported that providers recommended BGM 4–5 times/day compared to those who reported that their providers recommended BGM less than 4 times/day. Adjustment for age and insulin regimen did not change these results (Table 3, top).

A_{1c} also varied significantly by the number of times caregivers reported that BGM was conducted (Table 3, bottom, $p < .001$). Youth with caregivers who reported their child conducted BGM at least 6 times a day had significantly lower A_{1c} values than those who reported BGM fewer than 6 times a day; those who performed BGM 4–5 times/day had significantly lower A_{1c} values than those who conducted BGM fewer than 4 times/day. Adjustment for age and insulin regimen did not change these results (Table 3).

We then compared caregivers' report of provider recommendations for BGM to their report of their child's frequency of BGM. While most of the caregivers (92%) reported being told by their children's providers to conduct BGM at least 4 times/day (4–5 times or 6+ times/day), only 78% reported checking that

frequently. Table 4 shows that adherence to recommended monitoring frequency was reported by 62% of the caregivers with 24% reporting they test less frequently than recommended and 14% reporting they test more frequently. Of 10–12 year olds, 92% met or exceeded provider recommendations versus 72% of 13–15 year olds and 61% of >16 year olds ($p < .001$). Those with private insurance reported meeting or exceeding provider recommendations more than those without (79% versus 65%, $p = .01$), and those who met A_{1c} goal reported meeting or exceeding provider recommendations more than those who did not (91% versus 68%, $p = 001$).

A quarter of the participants test less frequently than recommended but 14% test more frequently than recommended.

In the adjusted analysis using multivariable logistic models, age (odds ratio [OR] = 8.6, 95% confidence interval [CI] = 3.8–19.6 for 10–12 year olds and OR = 1.4 (CI = 0.8–2.5) for 13–15 year olds, relative to > 16 year olds), private insurance (OR = 2.8, CI: 1.3–6.2), and A_{1c} Exceeded = 8.6 ± 1.4 , Met = 8.5 ± 1.6 , Did not meet = 9.7 ± 1.9 ; $p < .001$) predicted meeting or exceeding provider recommendations. Gender, family income, household composition, duration of diabetes, insulin regimen or depression scores were not significantly associated with reported adherence to recommended monitoring frequency.

These findings demonstrate that, among a large diverse sample of children with type 1 diabetes, most caregivers (92%) reported being told by their child's health care provider to conduct BGM at least

4 times/day. However, only 78% reported that their child monitored that often. Children whose caregivers reported that they monitored more frequently had lower A_{1c} values. Although causality cannot be determined from these cross-sectional data, previous reports have shown the detrimental effect of poor adherence to BGM monitoring recommendations on A_{1c} .³ Specifically, increased adherence has been associated with lower A_{1c} .^{1,9} Thus, our results reinforce the notion that BGM is strongly associated with glycemic control in children with type 1 diabetes.

One previous study reported that increasing BGM frequency to greater than 5 times a day did not result in further improvement of A_{1c} .¹⁰ In contrast, our findings suggest that those who reported checking at least 6 times/day did better than those who checked 4–5 times/day or less. Given this finding, it is interesting that only 22% of caregivers reported their provider recommended to check BGM 6 or more times per day. While frequency of provider recommendations did vary across study sites, no site differences were found in reported behaviors. In fact, compared to the percentage of caregivers reporting their provider recommended BGM >6 times/day, a slightly larger percentage of caregivers (26%) reported their child actually monitored that often. More attention to the accuracy of the 'topping out' factor cited by Ziegler et al.'s study and factors that may be involved in checking more than prescribed would be valuable, and may also have insurance policy coverage implications for BGM supplies.¹⁰

Age, race/ethnicity, and family variables including income and household structure emerged as important factors associated with reported recommendations for BGM, frequency of BGM, and adherence to provider recommendations.

Differences in age were particularly noteworthy as there was a significant difference in reported adherence to provider recommendations between the youngest and oldest children: only 60% of 16–17 year olds met or exceeded provider recommendations for BGM, compared with 93% of 10–12 year olds. These results are consistent with other studies which have demonstrated that older children with diabetes have poorer adherence to diabetes treatment than younger children.^{18,19}

Although results are consistent with other studies there are limitations to what can be concluded.

We also found that NHW participants reported being told to BGM more frequently than other race categories. While this study is not able to determine whether this is based on caregivers' perceptions or actual provider behavior, this finding may support growing literature showing racial disparities in diabetes outcomes,²⁰ and specifically corroborates reports indicating a disparity in regimen intensity for minority youth with type 1 diabetes.²¹ In addition, those on an insulin pump were more likely to report more frequent monitoring. This supports the findings of Cortina et al.,²² who also found that insulin pump users reported higher BGM than those on multiple daily injection regimens. Clearly there is a need to better understand the clinical implications, potential selection bias, and disparities that may exist surrounding intensity of the BGM recommendations children receive as well as their adherence behaviors.

There were several limitations to this study. Causality cannot be established from cross-sectional analyses.

Participation in the follow-up visit was sub-optimal, which may compromise the generalizability of these findings; for example, previous findings from the SEARCH study found that older children were less likely to participate in SEARCH study visits than younger children.²³ However, of the participants, response rate to the survey was high (94%). Further, the provider recommendations assessed and the actual frequency of BGM were both based on the report from the children's caregiver which was the parent or guardian who accompanied them to the study visit. We do not have their health care providers' report of their actual recommendations nor a confirmation of the actual frequency of BGM from either the children with diabetes or from downloading their meter as these measures were not collected as part of the study protocol. We therefore cannot be sure whether these reports, particularly those for the frequency of BGM, may accurately reflect their actual frequency of BGM, particularly among older teens who are more likely to be responsible for their own care. However, caregivers' recall or understanding of these recommendations, in particular, may be as important as what was actually recommended. It is reassuring to know that most caregivers report that health care providers recommended BGM at least 4 times/day. It is acknowledged that the caregiver report of actual BGM frequency may be overestimated. However, given the discordance reported between the provider recommendations and actual BGM frequency and the significant associations found with A_{1c} , we can be somewhat confident that these results, if inaccurate, may actually be conservative. Strengths of the study include sample size and the ethnic and geographic diversity of the children in the SEARCH study. In addition, measurement of A_{1c} was consistent

across centers and over time through the use of a single laboratory.

The need for age-specific interventions which focus on improving diabetes self-management is well-established.^{24,25} Understanding the demographic and clinical characteristics of children associated with adherence to BGM may elucidate the groups most at-risk for poor management and subsequent poor outcomes later in life. Targeting interventions to groups identified in this study who reported not knowing, or not following, these recommendations may be useful for clinicians aiming at improving glycemic outcomes in children with type 1 diabetes.



<http://www.diabetesresearchclinicalpractice.com>

Joyce P. Yi-Frazier

Joyce P. Yi-Frazier is a Research Assistant Professor at the University of Washington School of Medicine, Department of Pediatrics, Seattle, USA.